

## Viewpoint

## Driving a Hard Bargain with Diamond Qubits

Simon C. Benjamin

*Centre for Quantum Technologies, National University of Singapore, Singapore and  
Department of Materials, University of Oxford, Parks Rd, Oxford OX1 3PH, United Kingdom*

Jason M. Smith

*Department of Materials, University of Oxford, Parks Rd, Oxford OX1 3PH, United Kingdom*

Published October 3, 2011

*The coupling between nitrogen vacancy centers can be dramatically improved through the use of electromagnetic driving fields.*

Subject Areas: **Quantum Information, Spintronics****A Viewpoint on:****Electron-Mediated Nuclear-Spin Interactions between Distant Nitrogen-Vacancy Centers**

A. Bermudez, F. Jelezko, M. B. Plenio, and A. Retzker

*Physical Review Letters* **107**, 150503 2011 – Published October 3, 2011

The ideal building block or qubit for a quantum computer needs to possess several conflicting properties: it should be under precise control and yet almost perfectly isolated from its environment; it should be accessible separately from other components and yet coupled to them. A certain kind of impurity that occurs in diamond, called a nitrogen vacancy (NV) center, seems almost ideal, except that we don't yet know how best to couple different sites. This is absolutely essential in order to achieve entanglement between the various qubits. Now, writing in *Physical Review Letters*, Alejandro Bermudez and colleagues at the University of Ulm, Germany, present a theoretical scheme to improve NV coupling that exploits the complex structure of the NV system, together with an external electromagnetic driving field [1]. Information is stored in a nuclear spin at each location, while their big brother electron spins enable site-to-site communication. This elegant idea may prove less difficult to implement than previous solutions. If so, it will overcome a major obstacle to diamond-based quantum information technology.

A NV center is a “color center,” which gives pink diamond its hue. The system forms when two different kinds of defect bind together on adjacent lattice sites: a nitrogen impurity and a vacancy (a missing carbon atom) (see Fig. 1, bottom right). The combined entity may trap an additional electron, becoming negatively charged. The six unbound electrons present (three from adjacent carbons, two from the nitrogen atom, plus the one that has been trapped) form a spin-triplet ground state with exquisite coherence properties, which can be both polarized and measured using straightforward optical methods, and manipulated by applied electric and magnetic fields. Moreover, the presence of the nitrogen

atom (which has a nuclear spin of 1) implies that each NV center must have at least one nuclear spin present. Nuclei are ideal for storage of quantum information, because of their long coherence times. NV centers also benefit from the extreme hardness of diamond, which corresponds to an unusual phonon spectrum and thus allows certain quantum properties to persist, even at room temperature. Thus the NV center appears exceptionally well-suited for quantum information handling, and the past few years have seen a number of landmark demonstrations, from two and three-qubit operations using neighboring electronic or nuclear spins [2–4] to entanglement between an electron spin and an optical photon [5].

How can we enable different NV centers to achieve entanglement with one another, and thus produce a scalable architecture with the potential to represent the thousands, or millions, of qubits required for general-purpose quantum computers? Various solutions have been proposed. In one family of ideas, it is the optical emission of the centers that is exploited [6]. By making a joint measurement on photons emitted from two sites, those sites can be projected into an entangled state. This solution would therefore involve an optical network of well-separated NV centers, which is attractive in terms of scalability and the freedom to control and measure each site separately. However, NV centers are far from ideal photon sources; only a fraction of emissions occur without an accompanying phonon, the absence of which is required to produce entanglement. Another recent idea [7, 8] is to use a chain of “dark” spins—those on substitutional nitrogen atoms that do not have an adjoining vacancy—to link NV centers over distances of micrometers. The spins associated with such atoms are not optically active (and hence “dark”) but can nevertheless suffice to form a wire




FIG. 1: Lower right: Structurally, the NV center consists of a pair of adjacent defects in the diamond lattice, namely, a nitrogen atom substituting for a carbon, together with a vacancy (a missing atom). Main figure: In abstract terms, each NV center is composed of at least one nuclear spin, and an electron complex with total spin 1. Bermudez *et al.* have determined that under a suitable driving field, the electron spins will constitute a channel by which the nuclei in neighboring NV centers can interact, thus achieving entanglement. (APS/S. Benjamin and J. Smith)

or “bus” between the active sites. But again, there are challenges: to date, no group has demonstrated the control needed to implant such a heterogeneous network.

The idea put forward by the Ulm team is simpler to express than these previous suggestions. Since the nuclear spin at each site (bearing a qubit) is able to “see” its local electron spin partner via the hyperfine interaction, it follows that if the electron spins at neighboring sites can interact sufficiently strongly, then the two nuclei will be able to communicate indirectly through their electron brothers. While the idea is simple, it turns out to be more complex than one might imagine, in fact, Bermudez *et al.* start by establishing that the electron-mediated interaction between nuclei at adjacent NV sites will be far too weak to be useful! They estimate that even quite closely neighboring sites, 10 nanometers (nm) apart, will have an effective nuclear-nuclear coupling strength of only 0.1 hertz—requiring several seconds to achieve a useful exchange of information. This is impractical, since nuclear spins suffer dephasing (losing any stored qubit) in less than a second.

Happily, the Ulm team has discovered a solution. Their analysis predicts that when the spins are resonantly driven by externally applied electromagnetic fields, the effective strength of the interaction increases dramatically. In essence, they introduce a new energy scale into the problem, replacing the effect of the crystal field

splitting (which acts to suppress the effective nuclear-nuclear coupling) with the Rabi frequencies of the driven spins—a parameter that is under experimental control. Remarkably, with a suitable choice for this parameter, the coupling between nuclei is enhanced a thousandfold, becoming entirely practicable as a channel to exchange quantum information. As an added bonus, the act of driving the spins serves to protect the quantum state from the decohering effects of the surroundings. The effect is equivalent to “dynamic decoupling,” in which a spin that is periodically inverted at a frequency faster than the local magnetic field fluctuations acquires an aggregate zero phase [9].

Bermudez *et al.* have confirmed that their scheme should be possible with experimental conditions similar to those already achieved, in particular, in the work of Neumann *et al.*[10]. However, there are of course remaining challenges. To use this method to generate quantum entanglement on the sort of scale that would be useful for a general-purpose quantum computer would require a large array of NV centers. A recent study reports that positioning of individual centers can be achieved with about 80 nm accuracy, using implantation of  $CN^-$  molecules through nanoapertures [11]. This is approaching “the ballpark” for the new scheme and the authors speculate that denser arrays will indeed be possible. Even so, questions remain as to how the individual addressing of NV centers in a large group will be possible. The use of magnetic field gradients to impart a spatial dependence of the Zeeman splitting may well be part of the solution, but is unlikely to be sufficient without further discrimination. Nevertheless, the challenges in developing diamond quantum computers are being overcome at an astonishing rate. Bermudez *et al.*’s idea for entangling two nuclear spins is a powerful and remarkably elegant new option.

## References

- [1] A. Bermudez, F. Jelezko, M. B. Plenio, and A. Retzker, *Phys. Rev. Lett.* **107**, 150503 (2011).
- [2] R. Hanson *et al.*, *Phys. Rev. Lett.* **97**, 087601 (2006).
- [3] L. Childress *et al.*, *Science* **314**, 281 (2006).
- [4] P. Neumann *et al.*, *Science* **320**, 1326 (2008).
- [5] E. Togan *et al.*, *Nature* **466**, 730 (2010).
- [6] S. C. Benjamin, B. W. Lovett, and J. M. Smith, *Laser Photon. Rev.* **3**, 556 (2009).
- [7] N. Y. Yao *et al.*, *Phys. Rev. Lett.* **106**, 040505 (2011).
- [8] N. Y. Yao *et al.*, arXiv:1012.2864v1.
- [9] G. de Lange *et al.*, *Science* **330**, 60 (2010).
- [10] P. Neumann *et al.*, *Nature Phys.* **6**, 249 (2010).
- [11] P. Spinicelli *et al.*, *New J. Phys.* **13**, 025014 (2011).

## About the Authors

### Simon C. Benjamin



Simon Benjamin is the Senior Fellow in Quantum Materials in the Department of Materials at the University of Oxford. He lectures and leads a team of theorists broadly interested in the implementation of quantum devices (please see [qunat.org](http://qunat.org)). He is a Visiting Associate Professor at the Centre for Quantum Technologies in Singapore, where he jointly leads a group studying distributed quantum computing and other novel architectures.

### Jason M. Smith



Jason Smith is a Lecturer in the Department of Materials at the University of Oxford, where he is also a tutor and Fellow at Mansfield College. He received his D.Phil. in physics at the Clarendon Laboratory in Oxford in 1997. His research group studies the physics and applications of single photon emitters including color centers in diamond and semiconductor nanocrystals ([users.ox.ac.uk/~png](http://users.ox.ac.uk/~png)).