

Viewpoint

Is there a true supersolid phase transition ?

Sébastien Balibar

Laboratoire de Physique Statistique de l'Ecole Normale Supérieure, associé aux Universités Paris 6 et Paris 7 et au CNRS,
24 rue Lhomond 75005 Paris, France

Published May 10, 2010

New measurements of the rigidity of solid helium show that the emergence of supersolidity is actually a crossover, rather than a true phase transition.

Subject Areas: **Superfluidity**

A Viewpoint on:

Frequency Dependence and Dissipation in the Dynamics of Solid Helium

Oleksandr Syshchenko, James Day and John Beamish

Phys. Rev. Lett. **104**, 195301 (2010) – Published May 10, 2010

Six years have passed since Kim and Chan [1, 2] discovered supersolidity when they found that solid ^4He does not rotate like a classical solid. Despite substantial effort, however, this phenomenon is not yet understood [3]. Some probably think that the research on this puzzling issue is too slow. Others, including myself, enjoy every new controversy and every new advance as if they were reading a detective story. In this tale, the latest twist is a study reported in *Physical Review Letters* by Oleksandr Syshchenko, James Day, and John Beamish at the University of Alberta, Canada [4], of the shear modulus of solid ^4He in the temperature range where it becomes supersolid. They confirm that the observed change in stiffness is associated with a binding of dislocations to ^3He impurities (Fig. 1) when the solid is cooled down (then unbinding when warming up, of course) and they show now that this binding is gradual because there is a rather wide distribution of binding energies. Why is this so important?

As is now well known, supersolidity means a coexistence of superfluid and solid behaviors. This is surprising but possible, especially if the solid under consideration has disorder, for example, dislocations [5]. Even more surprising was probably the discovery by Day and Beamish [6] in 2007 that solid helium is stiffer in its supersolid phase than in its normal phase. Moreover, the variation of the elastic shear modulus as a function of temperature was shown to be exactly the same as that of the rotational inertia. The two properties must be consequences of a single phenomenon, but which one? This is what all of the researchers in this field are trying to understand.

Day and Beamish had also shown that, near the transition, the elastic modulus has a much larger variation than the rotational inertia. The typical variation of the modulus is 10% in polycrystalline samples instead of only 1% for the inertia. Now, Syshchenko *et al.*[4] show




FIG. 1: According to Syshchenko *et al.*[4], the binding of dislocations to ^3He impurities is the mechanism that controls the change in shear modulus of solid helium at the temperature where it becomes supersolid. With mobile dislocations (left panel), the crystal is softer than if dislocations are pinned by ^3He impurities (right panel). Syshchenko *et al.* prove that the transition is a broad crossover because the distribution of binding energies is very wide. But the mobility of dislocation lines is due to the existence of kinks, which are mobile along the lines. As a consequence, one interesting question remains: Is it possible that, even in the absence of ^3He impurities, the dislocations become smooth and fixed straight lines without kinks at low temperature, due to the periodic potential of the crystal lattice? This is what Aleinikava *et al.*[10] proposed and this now needs to be checked. (Illustration: Adapted from [3])

that the elastic properties can also be studied in a wide range of frequencies—from 0.5 Hz to 8 kHz—while the study of the rotational inertia in torsional oscillators is restricted to a few hundred hertz around 1 kHz. They took advantage of that to study the nature of the transition from the normal to the supersolid state. Some authors had tried to fit variations as a function of temperature with power laws and critical exponents, as is usually done close to phase transition temperatures [7]. But this work shows that the latter approach is not appropriate. The onset of the transition is broad because

the binding energies of dislocations to ^3He impurities are distributed around a temperature of 0.73 K with a large width (0.45 K). It is a crossover in a thermally activated process and, according to Syshchenko *et al.*: “if there is a true phase transition ... it must be at lower temperature.”

Is there a “true phase transition” then? Syshchenko, Day, and Beamish probably have their own thoughts on the matter, but they are careful and do not answer this fundamental question. However, I will attempt an answer of my own. One needs to understand first why there is a coupling between rigidity and supersolidity. The reason why helium crystals are stiffer at low temperature, say 20 mK, than at higher temperature, say 200 mK, is that a crystal with fixed dislocations is stiffer than a crystal with mobile dislocations. This is actually well known in materials science [8]. So I propose the following scenario: Inside the core of dislocations, the atoms are not densely packed, and that is actually why atom exchange is much easier there than in the bulk of the crystal, something which may lead to quantum coherence and superfluidity. But if the dislocations are mobile, they have transverse fluctuations, that is, fluctuations of their position and consequently of their total length if their ends are fixed. As a result, there should be fluctuating mass currents. In a quantum coherent system, mass currents are proportional to the gradient of the phase of the wave function so that these fluctuations should induce phase fluctuations and this should kill the quantum coherence that is necessary for supersolidity.

My intuition is that a fluctuating dislocation cannot be supersolid. On the contrary, if a dislocation is supersolid, the phase of the wave function must be locked and the dislocation a fixed line. This is how I understand that the stiffness transition in helium crystals is closely linked to their supersolid transition. When the dislocations are pinned by ^3He impurities, that is, at low temperature, they form a network of supersolid lines throughout the whole solid, but if the dislocations unpin, they fluctuate and supersolidity disappears. If my physical intuition is correct, the dislocation pinning controls the whole phenomenon. In this picture, the onset of the phenomenon may be broad, as observed by Syshchenko *et al.*, but as the temperature is lowered, all dislocations progressively stop fluctuating and a quantum coherence should set in and percolate through the whole network of dislocations, ending with a “true phase transition” to a supersolid state. This must be the origin of the peak in the specific heat that has been measured by Lin *et al.*[9]. Now how to check this picture?

Last year, my colleagues and I had shown that it is possible to suppress all ^3He impurities [10], but we


doubt that this is sufficient to free all dislocations at all temperatures and suppress supersolidity. Indeed, we have also found that, in ultrapure samples, the “crossover” to a stiff state—I do not call it a transition anymore—is displaced to lower temperature (less than 40 mK) but perhaps not suppressed [11]. One possible interpretation of this can be found in an article by Aleinikava *et al.*[12] who explained in 2010 that the dislocations could be pinned not only by ^3He impurities but also by the crystal lattice itself (its so-called “Peierls potential”). There should be a roughening transition of the dislocations lines, which should depend on the presence of impurities but is not suppressed in their absence. Given all these other results and in order to check the dislocation scenario for supersolidity, the thing to do is to suppress not only impurities but also dislocations in single crystals and see if supersolidity disappears. It should—except if a totally different scenario is at work, for example, the proliferation of vortices, which is proposed by Anderson [13] and was recently presented as supporting their study of relaxation times by Kim *et al.*[14]. We will see. I actually expect another controversy to come up as a result of a very recent article by Reppy [15] in which he presents measurements in a highly disordered sample that are incompatible with supersolidity. In view of this, I am afraid that our detective story is far from finished.

References

- [1] E. Kim and M. H. W. Chan, *Nature* **427**, 225 (2004).
- [2] E. Kim and M. H. W. Chan, *Science* **305**, 1941 (2004).
- [3] For a recent review and see S. Balibar, *Nature* **464**, 176 (2010).
- [4] O. Syshchenko, J. Day, and J. Beamish, *Phys. Rev. Lett.* **104**, 195301 (2010).
- [5] M. Boninsegni, A. B. Kuklov, L. Pollet, N. V. Prokof'ev, B. V. Svistunov, and M. Troyer, *Phys. Rev. Lett.* **99**, 035301 (2007).
- [6] J. Day and J. Beamish, *Nature* **450**, 853 (2007).
- [7] A. C. Clark, J. West, and M. H. W. Chan, *Phys. Rev. Lett.* **99**, 135302 (2007).
- [8] J. Friedel, *Dislocations* (Pergamon Press, Oxford, 1964).
- [9] X. Lin, A. C. Clark, and M. H. W. Chan, *Nature* **449**, 1025 (2007); X. Lin, A. C. Clark, Z. G. Cheng, and M. H. W. Chan, *Phys. Rev. Lett.* **102**, 125302 (2009).
- [10] C. Pantalei, X. Rojas, D. O. Edwards, H. J. Maris, and S. Balibar, *J. Low Temp. Phys.* **159**, 452 (2010).
- [11] X. Rojas, C. Pantalei, H. J. Maris, and S. Balibar, *J. Low Temp. Phys.* **158**, 478 (2010).
- [12] D. Aleinikava, E. Dedits, A. B. Kuklov, and D. Schmeltzer, *Europhys. Lett.* **89**, 46002 (2010).
- [13] P. W. Anderson, *Nature Phys.* **3**, 160 (2007).
- [14] H. Choi, S. Kwon, D. Y. Kim, and E. Kim, *Nature Phys.* **6**, 4 April (2010).
- [15] J. D. Reppy, arXiv:1004.1129.

About the Author

Sébastien Balibar



Sébastien Balibar is a CNRS Director of Research working at the Laboratoire de Physique Statistique de l'Ecole Normale Supérieure (ENS), Paris. He received his Ph.D. in 1976 under Albert Libchaber at the ENS and has carried out extensive research in low-temperature and condensed-matter physics. Dr. Balibar was an invited professor at Konstanz Universität, Kyoto University, and Harvard University, where he was a recipient of the Loeb Lecture-ship. Among various honors, he was elected a fellow of the American Physical Society and shared the Fritz London Memorial Prize in 2005. The author of numerous research papers, he has also written the book *La Pomme et l'Atome*, (Odile Jacob, Paris, 2005) (*The Atom and the Apple: Twelve Tales of Contemporary Physics*, Princeton University Press, 2008).