

Viewpoint

Free-Electron Lasers Trigger Nuclear Transitions

Félicie Albert

Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

Published February 24, 2014

New theoretical calculations show that an x-ray free-electron-laser pulse can generate transitions in excited nuclei via an indirect process involving electron capture.

Subject Areas: **Nuclear Physics**

A Viewpoint on:

Dominant Secondary Nuclear Photoexcitation with the X-Ray Free-Electron Laser

Jonas Gunst, Yuri A. Litvinov, Christoph H. Keitel, and Adriana Pálffy

Physical Review Letters **112**, 082501 2014 – Published February 24, 2014

Since they became available to users in 2009 [1], hard x-ray free-electron lasers (XFELs) have been transformational tools for many scientific disciplines. These kilometer-sized machines generate x rays with unprecedented brightness at energies above 1 kilo-electron-volt and in pulses as short as a few femtoseconds (10^{-15} s). It is not always entirely clear how these bright pulses interact with targets. In the case of nuclear physics, the expectation was that XFEL light would predominantly interact with nuclei through direct photoexcitation. But a new theoretical study of excited nuclear states has revealed that, surprisingly, XFEL-nuclear interactions are dominated by a secondary mechanism. As described in *Physical Review Letters* [2], this indirect photoexcitation mechanism, which involves electron capture from the laser-induced plasma, is over a million times more efficient. The results may be interesting for future energy storage applications, in which excited nuclei perform as nuclear batteries.

Free-electron lasers are a big step up in x-ray generation, delivering 10^8 more photons per volume than synchrotron sources. The remarkable performance of XFELs is achieved by accelerating high-quality, ultrashort electron bunches to giga-electron-volt energies in a linear particle accelerator. Radiation is emitted from the periodic transverse oscillations of the electron bunch passing through an undulator—a periodic arrangement of magnets. The radiation interacts with the electron bunch, which yields density modulations on the bunch. This density modulation enhances the x-ray emission process, and lasing can occur. There are currently three operational XFELs in the world: the LCLS in the US, SACLA in Japan, and FLASH in Germany, with a prolific scientific production and ever-increasing experimental demands and number of users. Seminal achievements include the creation of some of the hottest plasmas on earth [3], the first atomic inner-shell x-ray laser [4], and structural studies of proteins with crystallography [5]. In these

examples, x rays play a role in understanding atomic and molecular processes.

While they hold the promise to revolutionize atomic physics and structural biology, XFELs could also help achieve breakthroughs in nuclear physics. Jonas Gunst of the Max Planck Institute for Nuclear Physics in Heidelberg, Germany, and his colleagues report the unexpected result that XFELs can indirectly trigger nuclear excitation processes in nuclear isomers by coupling nuclei to the atomic shell [2]. Nuclear “isomers” [6], a term borrowed from chemistry, are excited states of a nucleus that have the same number of neutrons and protons (as opposed to isotopes, having a different number of neutrons). Typical half-lives of isomeric states are on the order of 1 picosecond, but some of them can last several hours. According to the findings of Gunst *et al.*, triggering these excited states to release their energy on demand seems within reach. The authors describe the interaction of an XFEL with one of the isomeric states of molybdenum, ^{93m}Mo , which has an excitation energy of 2.5 mega-electron-volts and a lifetime of nearly 7 hours. Nuclear decay of the isomeric level can be enhanced by exciting the nucleus to a less stable, intermediate level lying above the isomer. X rays from the XFEL can perform this transition via two different mechanisms.

The first route is the direct one: if the XFEL is tuned to 4.85-kilo-electron-volts, then the x rays can excite the nuclei resonantly to a higher excitation state, from which they rapidly decay back to the ground state, releasing x rays and gamma rays. This direct photoexcitation process is, however, rather inefficient because of the fact that the energy width of the nuclear transition is 8 orders of magnitude smaller than the width of the XFEL spectrum. Therefore, only a small fraction of the laser photons are resonant with the nuclear transition.

The second possible route for releasing the isomer’s energy is indirect, so it was thought to be less important than the direct one. This secondary mechanism,

called nuclear excitation by electron capture, or NEEC, is a resonant process, in which the XFEL light removes electrons from the target atoms (in the isomeric excited state), and these free electrons form a plasma with a thermal energy of about a few hundred electron volts (see Fig. 1). An electron from this plasma can subsequently be captured into an empty state of the atomic shell (left vacant from the interaction of the XFEL with the atom). This captured electron gives its excess energy to excite a nuclear state. First described in 1976 [7], the NEEC process is considered as a possible nuclear decay mechanism in stars, but previous experiments with synchrotron or bremsstrahlung radiation have not had the required photon density or energy for creating the phenomenon in the lab. Presumably, the light from one of the new XFEL facilities will be able to ionize enough electrons from the target to create the plasma and bring it to a high temperature.

Assuming an XFEL source of photons, Gunst *et al.* calculate that indirect photoexcitation through the NEEC process is at least 6 orders of magnitude more efficient than direct photoexcitation. One reason for this large discrepancy is that—unlike the direct photoexcitation of the 4.85 kilo-electron-volt transition—the NEEC mechanism is less sensitive to the resonance condition, since the photons have a wide range of inner-shell electrons with which to interact. Moreover, the plasma lasts much longer than the XFEL pulse, so there is more time for electron capture to occur as opposed to direct photon-mediated excitation.

The consequences of this work could spark new life into the field of nuclear energy storage, where scientists have been studying for years the prospects of using metastable nuclear isomers in high-energy batteries. The advantage of isomers, which are typically produced in nuclear reactions or through irradiation, is that they provide nuclear power with no long-lasting nuclear waste. However, the challenge is the control of the release of the stored nuclear energy. Gunst *et al.* have shown that the light from an XFEL could trigger the rapid decay of a nuclear isomer. Now, given that this process requires using a large and costly facility, it seems unlikely that there will be a practical energy-storage device in the near future, but this new work could stimulate research on other triggering schemes. Another application could be in the medical community. Long-lived nuclear isomers are of interest for a number of applications in nuclear medicine [8], where they can be used for therapeutic (endotherapy) or diagnostic (nuclear imaging) purposes. In the example of ^{93m}Mo triggering described by Gunst *et al.*, a 1-mega-electron-volt photon is emitted in the decay cascade from the triggering level to the ground state, and it can potentially be detected by modern imaging systems or target a specific tumor.

The authors are marching into a virtually virgin territory, as no existing x-ray source has yet enabled the experimental demonstration of NEEC triggering of isomer relaxation. This theoretical work should inspire ex-




FIG. 1: This graphic shows the secondary process by which XFEL light can drive the energy release from a sample containing a nuclear isomer. In the first step (top), photons knock electrons out of inner atomic shells, leading to a plasma of free electrons and ionized atoms. This interaction does not directly affect the nucleus, as it remains in the metastable isomeric state (top right). However, when electrons from the plasma are later captured in the inner-shell vacancies, their excess energy can excite the nucleus to a higher, but less stable, energy state (middle). In the last step, the nucleus relaxes to its ground state, releasing photons in a cascade (bottom). (APS/Alan Stonebraker)

perimental efforts at XFEL facilities that have sufficient photon energy and x-ray intensity to allow the NEEC process to occur. Researchers can isolate the NEEC signal by tuning their XFEL light out of resonance with direct photoexcitation, so that the only isomer emission should be from the indirect mechanism. The effort will be challenging, but the results could be a major breakthrough.

References

- [1] P. Emma *et al.*, “First Lasing and Operation of an Angstrom-Wavelength Free-Electron Laser,” *Nature Photon.* **4**, 641 (2010).
- [2] Jonas Gunst, Yuri A. Litvinov, Christoph H. Keitel, and Adriana Pálffy, “Dominant Secondary Nuclear Photoexcitation with the X-Ray Free-Electron Laser,” *Phys. Rev. Lett.* **112**, 082501 (2014).
- [3] S. M. Vinko *et al.*, “Creation and Diagnosis of a Solid-Density Plasma with an X-ray Free-Electron laser,” *Nature* **482**, 59 (2012).
- [4] N. Rohringer *et al.*, “Atomic Inner-Shell X-Ray Laser at 1.46 Nanometres Pumped by an X-Ray Free-Electron laser,” *Nature* **481**, 488 (2012).
- [5] H. N. Chapman, P. Fromme, A. Barty, T. A. White, R. A. Kirian, A. Aquila, and M. S. Hunter, “Femtosecond X-Ray Protein Nanocrystallography,” *Nature* **470**, 73 (2011).
- [6] P. M. Walker and J. J. Carroll, “Ups and Downs of Nuclear Isomers,” *Phys. Today* **58**, No. 6, 39 (2005).
- [7] V. I. Goldanskii and V. A. Namiot, “On the Excitation of Isomeric Nuclear Levels by Laser Radiation through Inverse Internal Electron Conversion,” *Phys. Lett. B* **62**, 393 (1976).
- [8] D. Habs and U. Koster, “Production of Medical Radioisotopes with High Specific Activity in Photonuclear Reactions with Γ -Beams of High Intensity and Large Brilliance,” *Appl. Phys. B* **103**, 501 (2011).

About the Author

Félicie Albert



Félicie Albert is an experimental plasma physicist at the National Ignition Facility, Lawrence Livermore National Laboratory, who earned her Ph.D. in physics from the Ecole Polytechnique in France in 2007. She has conducted many experiments using high-intensity lasers in France and the US. Her areas of interest include the generation of novel sources of electrons, x rays, and gamma rays through laser-plasma interaction, Compton scattering, and laser-wakefield acceleration. She is currently leading an effort to develop these sources for applications in high-energy density sciences as well as doing experiments for the National Ignition Facility.