

Rapid Neutrino Cooling in the Neutron Star MXB 1659-29Edward F. Brown,^{1,*} Andrew Cumming,^{2,†} Farrukh J. Fattoyev,^{3,‡}C. J. Horowitz,^{3,§} Dany Page,^{4,||} and Sanjay Reddy^{5,¶}¹*Department of Physics and Astronomy, Michigan State University, 567 Wilson Road, East Lansing, Michigan 48864, USA*²*Department of Physics and McGill Space Institute, McGill University, 3600 rue University, Montreal, Quebec, Canada H3A 2T8*³*Center for Exploration of Energy and Matter and Department of Physics, Indiana University, Bloomington, Indiana 47405, USA*⁴*Instituto de Astronomía, Universidad Nacional Autónoma de México, México, CDMX 04510, Mexico*⁵*Institute for Nuclear Theory, University of Washington, Seattle, Washington 98195, USA* (Received 20 July 2017; revised manuscript received 20 December 2017; published 30 April 2018)

We show that the neutron star in the transient system MXB 1659-29 has a core neutrino luminosity that substantially exceeds that of the modified Urca reactions (i.e., $n + n \rightarrow n + p + e^- + \bar{\nu}_e$ and inverse) and is consistent with the direct Urca ($n \rightarrow p + e^- + \bar{\nu}_e$ and inverse) reaction occurring in a small fraction of the core. Observations of the thermal relaxation of the neutron star crust following 2.5 yr of accretion allow us to measure the energy deposited into the core during accretion, which is then reradiated as neutrinos, and infer the core temperature. For a nucleonic core, this requires that the nucleons are unpaired and that the proton fraction exceeds a critical value to allow the direct Urca reaction to proceed. The neutron star in MXB 1659-29 is the first with a firmly detected thermal component in its x-ray spectrum that needs a fast neutrino-cooling process. Measurements of the temperature variation of the neutron star core during quiescence would place an upper limit on the core specific heat and serve as a check on the fraction of the neutron star core in which nucleons are unpaired.

DOI: [10.1103/PhysRevLett.120.182701](https://doi.org/10.1103/PhysRevLett.120.182701)

The composition and phases of matter at a low temperature and supranuclear density are fundamentally important to our understanding of QCD under extreme conditions and in astrophysics. Such matter is found in the cores of neutron stars, and observations of the surface emission of neutron stars allow us to infer their internal temperatures and hence the efficiency of neutrino cooling in their interiors. The neutrino emissivity depends on the composition of the core and whether the constituent particles are paired, forming a superfluid state [1,2]. Therefore, observations showing that rapid neutrino cooling is operating would have important implications for our understanding of dense matter [3,4].

It is useful to classify neutrino emissivities based on their temperature dependence. The first is fast, or direct Urca-like reactions ($n \rightarrow p + e^- + \bar{\nu}_e$ and inverse), for which the emissivity $\epsilon_{\text{fast}} = Q_f T_8^6$. Here we use the notation $T_8 \equiv T/10^8$ K. The coefficient Q_f is a weak function of the baryon density but depends on the phase structure: $Q_f \simeq 10^{35}\text{--}10^{36}$ erg km⁻³ s⁻¹ for nucleon direct Urca in normal nuclear matter [5], $Q_f \simeq 10^{32}\text{--}10^{35}$ erg km⁻³ s⁻¹ for nucleon direct Urca in the presence of a meson condensed phase [6,7], and $Q_f \simeq 10^{34}\text{--}10^{36}$ erg km⁻³ s⁻¹ for direct Urca reactions in quark matter [8,9]. The second class, intermediate in strength, are Cooper pair breaking and formation (PBF) processes, for which $\epsilon_{\text{int}} = Q_i T_8^7$ and

$Q_i \simeq 10^{28}\text{--}10^{29}$ erg km⁻³ s⁻¹ $\times F_{\text{PBF}}(T_c/T)$. This intermediate process operates only in regions where $T \lesssim T_c$, T_c being the critical temperature for neutron superfluidity, since $F_{\text{PBF}}(T_c/T) \rightarrow 0$ otherwise [10–12]. The third class are slow, or modified Urca-like, reactions [13], for which $\epsilon_{\text{slow}} = Q_s T_8^8$ and $Q_s \simeq 10^{28}\text{--}10^{29}$ erg km⁻³ s⁻¹ is again a weak function of the baryon density.

It is not known if direct Urca occurs in neutron stars, because the reactions are blocked by momentum conservation unless the proton fraction exceeds a critical value, $Y_p = 0.11\text{--}0.15$ [5]. Furthermore, the pairing of protons or neutrons in the core would suppress the direct Urca reactions, so that the neutrino luminosity emitted by the direct Urca process also depends on the density dependence of the pairing energy. The classical test for the presence of enhanced cooling is to observe a sample of isolated cooling neutron stars (for a catalog, see Ref. [14] and references therein) and compare with theoretical cooling models (for a review, see [1,2]). Of these, some may need enhanced emission [15]. Accreting neutron stars in low-mass x-ray binaries have been used to place constraints on the neutrino emissivity of the neutron star core [16]. For transient accretors, the neutron star core comes into thermal balance between accretion-induced heating and neutrino cooling. During quiescence, when the accretion halts, the surface temperature can be measured and used to infer the core

temperature. For two systems, SAX J1808.4-3658 and 1H 1905, there are only upper limits on the thermal emission from the neutron star in quiescence [17,18]. These upper limits are stringent enough to suggest that fast neutrino cooling is operating, but the thermal emission from these stars has not been directly detected.

The low-mass x-ray binary MXB 1659-29 went into quiescence in 2001 after a 2.5 yr outburst [19]. Observations with *Chandra* and *XMM* over the next 15 yr showed a declining neutron star surface temperature [20–23]. MXB 1659-29 had previously been in an outburst 21 yr earlier (see [21] and references therein). In 2015, a new outburst was reported [24], providing a second chance to follow the cooling of the source when it goes back into quiescence. Here we show that if the outburst-quiescent cycles observed to date represent the long-term average (over the thermal timescale of the core [25]) accretion behavior of the source, then the core neutrino luminosity must be a fast cooling process consistent with direct Urca reactions occurring in $\sim 1\%$ of the neutron star core.

Before presenting detailed numerical calculations, we first give a basic argument. Nuclear reactions in the crust (where the density is below saturation) during accretion deposit (see, e.g., [26]) ≈ 2 MeV per accreted nucleon (about 2×10^{18} erg g $^{-1}$). Over the 2.5 yr outburst of MXB 1659-29, the neutron star accumulated a mass of $\approx 8 \times 10^{24}$ g; as a result, the total energy deposited by these crust reactions during the outburst is $E \approx 2 \times 10^{43}$ erg. Most of this heat is conducted into the core. If the core is in a long-term thermal equilibrium—established over the previous ≈ 3 –30 outbursts for the cold core of this source—this deposited energy is radiated by neutrinos between outbursts, giving a neutrino luminosity of 2×10^{43} erg/20 yr $\approx 3 \times 10^{34}$ erg s $^{-1}$. With knowledge of the core temperature, we can use this luminosity to constrain the neutrino emission process.

Reference [27] used the observed effective temperature, after the crust had thermally relaxed, of 55 eV [28] to derive a redshifted core temperature $\tilde{T} = 2.5 \times 10^7$ K. (The redshifted core temperature, denoted by \tilde{T} , is the temperature measured by an observer at infinity and is the quantity that is uniform over an isothermal core.) For fast neutrino cooling, $L_\nu \propto \tilde{T}^6$, we infer that

$$L_\nu \approx 10^{38} \text{ erg s}^{-1} \tilde{T}_8^6. \quad (1)$$

The temperature in a local frame is $T = \tilde{T}e^{-\phi}$, where $\phi \equiv \frac{1}{2} \ln g_{00} \approx -0.5$ is the gravitational potential found by integrating the Tolman-Oppenheimer-Volkoff equation (for details, see [27]). We compare Eq. (1) with the integral of the emissivity for direct Urca, $\epsilon_{\nu,\text{dU}} \sim 10^{36}$ erg km $^{-3}$ s $^{-1}$ T $_8^6$, over a volume V_{dU} . For direct Urca reactions, having only the innermost ≈ 3 km ($\lesssim 2\%$ of the volume of the core) of the neutron star above threshold is sufficient to supply this

neutrino luminosity. Conversely, an emission process that is $\lesssim 10^{-2}$ as strong as direct Urca cannot supply the required neutrino luminosity, even if it were acting over the whole core volume.

Since the inferred core temperature is small, slow reactions with rates $\propto T^8$, such as the modified Urca, $nn \rightarrow npe^- \bar{\nu}_e$, $np \rightarrow ppe^- \bar{\nu}_e$, and their inverses, are greatly suppressed, relative to the direct Urca rate, by a factor of $\approx 10^{-1} (m_n/m_\pi)^4 (k_B \tilde{T}/m_n c^2)^2 \simeq 10^{-7}$ at $\tilde{T} = 10^8$ K. Here m_n and m_π denote the neutron and pion masses, respectively. Similarly, even if relatively large regions of the star were able to sustain the intermediate class of PBF neutrino reactions with $L_{\nu,\text{PBF}} \propto \tilde{T}^7$, the integrated neutrino luminosities would be too small. This large difference between neutrino luminosities at the low inferred core temperature allows us to unambiguously identify the occurrence of rapid cooling in MXB 1659-29.

To confirm this estimate, we calculated a time-dependent model of the response of the core to many accretion outbursts. Our calculation follows the crust physics in detail [29] but represents the core as a single zone with a total heat capacity and neutrino luminosity. This is reasonable as the thermal diffusion timescale of the core is $\ll 1$ yr [rescaling Eq. (16) of Ref. [27] to $\tilde{T}_8 = 0.25$]. To find the long-term steady state, we started with a guess for the core temperature just before the first outburst, ran a sequence of 50 outbursts (~ 1000 yr), and allowed the core to come into equilibrium. Once the equilibrium core temperature was determined, we reran the sequence of outbursts with the equilibrium core temperature as a starting point. We repeated this sequence, adjusting other parameters such as the crust impurity parameter to fit the observed cooling curve for MXB 1659-29. Although we used a uniform outburst accretion rate and identical outburst-recurrence durations, we find that our conclusions are not sensitive to this choice (cf. Ref. [27]). The results are shown in Fig. 1. We find that a core neutrino luminosity of 2.1×10^{38} erg s $^{-1} \tilde{T}_8^6$ provides a good fit (luminosities are given as measured by an observer at infinity).

To assess the range of allowed values for L_ν/\tilde{T}_8^6 , we fit the observed x-ray emission using a Markov chain Monte Carlo (MCMC) algorithm. For these runs, we calculated only one outburst and kept the core temperature fixed during the simulation. By measuring the total energy conducted into the core during the outburst and quiescence, we then calculated the neutrino luminosity required for the core to be in equilibrium at the assumed core temperature. To test the robustness of our inference about L_ν/\tilde{T}_8^6 , we calculated three cases with different assumed priors. The resulting posterior distributions of L_ν/\tilde{T}_8^6 are shown in Fig. 2. The first uses a prior for the impurity parameter Q_{imp} (a higher Q_{imp} implies a lower thermal conductivity, and hence a longer thermal relaxation, of the crust) that is uniform in $\log_{10} Q_{\text{imp}}$ (as was done in Ref. [27]). Because




FIG. 1. An example model in long-term equilibrium with core neutrino emissivity $L_\nu = 2.1 \times 10^{38} \text{ erg s}^{-1} \tilde{T}_8^6$ and core heat capacity $C = 10^{37} \text{ erg K}^{-1} \tilde{T}_8$. We assume identical 2.5 yr outbursts and that the source spends 5500 days (≈ 15 yr) in quiescence. The observations are indicated as black points; in the last observation (gray point), the count rate decreased, which may be due either to increased absorption or to cooling. We assume that the neutron star temperature remains unchanged in our fits. Other parameters are $M = 1.6 M_\odot$, $R = 11.2 \text{ km}$, and $Q_{\text{imp}} = 3.3$. After ≈ 2000 days into quiescence, when the crust has come into thermal equilibrium with the core, the effective temperature continues to decline as the core cools, at a rate $\Delta T_{\text{eff}}^\infty / T_{\text{eff}}^\infty \approx 6.5\%$ per decade.

this gives much greater weight to $Q_{\text{imp}} \ll 1$, low values of surface gravity are preferred. While massive neutron stars (as needed to achieve the high densities required for direct Urca reactions) are allowed, they also need to have large radii to produce a low surface gravity. If instead we use a prior that is uniform in Q_{imp} , the range of surface gravity broadens and allows for more compact massive neutron stars. For the uniform prior on Q_{imp} , we find a best-fit value of $\log_{10} L_\nu / \tilde{T}_8^6 = 38.2 \pm 0.22$, corresponding to a central value of $1.6 \times 10^{38} \text{ erg s}^{-1}$ with a 1-sigma standard deviation of a factor of 2 in each direction. In Fig. 2, the green histogram shows what happens when we impose a lower limit of $2 M_\odot$ on the mass. In all cases, the preferred neutrino luminosity is $L_\nu / \tilde{T}_8^6 \sim 10^{38} \text{ erg s}^{-1}$.

The last observation, approximately 4000 days after the end of the outburst, had a decrease in the x-ray count rate. There are two possible explanations, with different implications for the inferred T_{eff}^∞ , as indicated by the two gray points in Fig. 1. The neutron star surface temperature may have cooled to a lower value of $< 50 \text{ eV}$ [23]. This could be due to relaxation of the nuclear pasta layer [30,31]. The observed decrease in the x-ray count rate can also be




FIG. 2. Posterior distribution of the neutrino cooling prefactor L_ν / \tilde{T}_8^6 [see Eq. (1)] from the MCMC fits to the MXB 1659-29 cooling curve. To illustrate the robustness to different prior assumptions, we show results for three different priors: uniform in $\log_{10} Q_{\text{imp}}$ (black curve), uniform in Q_{imp} (blue curve), and uniform prior in Q_{imp} with the further restriction that $M > 2 M_\odot$ (green curve).

explained by an increased x-ray absorption from matter accumulating in a disk within the binary (see the discussion in Ref. [23]). In the fits above, we assume that the neutron star temperature remained unchanged at 55 eV (upper gray point). A lower core temperature of 50 eV (lower gray point) would increase our inferred cooling rate L / \tilde{T}_8^6 by approximately a factor of 2.

The derivation of the core temperature from the observed effective temperature depends on the composition assumed for the neutron star envelope. We have assumed a light element (He) envelope, giving a core temperature of $\tilde{T} = 2.5 \times 10^7 \text{ K}$ for the observed effective temperature of 55 eV. A heavy element (Fe) envelope gives a larger core temperature $\tilde{T} = 5.5 \times 10^7 \text{ K}$ [Eq. (5) in Ref. [27]], reducing the inferred value of L_ν / \tilde{T}_8^6 by a factor of ~ 100 . However, as discussed in Ref. [27], the envelope composition changes the shape of the cooling curve. In the case of an Fe envelope, the hotter crust gives a much smoother decline in temperature (Fig. 1 in Ref. [27]). By varying other parameters, we can find acceptable fits to the data with an Fe envelope but only by increasing the accretion rate (and therefore deep crustal heating rate) by a factor of 5–10, which would be inconsistent with the observed persistent x-ray luminosity during the outburst. In addition, a light element envelope accretes quickly ($\sim 10^4 \text{ s}$ at typical mass accretion rates) and is therefore more likely to be left over at the end of the accretion outburst. We therefore use a light element envelope in our fits. A heavy element envelope was used in Ref. [32], who placed MXB 1659-29 in the diagram of quiescent luminosity against a time-averaged accretion rate. Although our use of a light element envelope does not change the location of the contours of neutrino emissivities by a substantial amount in this

diagram ($\lesssim 2$; see Ref. [1]), it has a significant effect on the inferred value of L_ν/\tilde{T}^6 .

The effective temperature 55 eV was obtained assuming that the distance to MXB 1659-29 is 10 kpc [28]. A 30% larger distance [33] gives T_{eff} larger by $\approx 7\%$ [22] and \tilde{T} larger by $\approx 15\%$. The resulting factor of 2–3 in \tilde{T}^6 is offset by an increased inferred accretion rate and heating rate for a larger distance, so that overall we expect distance uncertainties to affect our inferred L_ν/\tilde{T}^6 by less than a factor of 2.

Monitoring observations of MXB 1659-29 when in quiescence, after its crust has thermally relaxed, can constrain the specific heat of the core. The cooling $\Delta T_{\text{eff}}^\infty/T_{\text{eff}}^\infty$ over some fixed time is $\propto L_\nu/C$ [27] (Fig. 3, dashed lines) for an inferred \tilde{T} and E . Figure 3 shows that any constraint on cooling after the next outburst will constrain the heat capacity of the core. For example, if the decrease in the effective temperature, after the crust has thermally relaxed, is $\lesssim 1\%$, then a large fraction of the nucleons in the core should be normal (upper horizontal gray band), with $C/\tilde{T}_8 \gtrsim 10^{38}$ erg K $^{-1}$. Conversely, if the core cools significantly ($\gtrsim 4\%$) during quiescence, this would suggest that most of the nucleons are paired and




FIG. 3. Constraints on possible values of the specific heat C and neutrino luminosity $L_\nu \propto \tilde{T}^6$ for the \tilde{T} evaluated at the best-fit core temperature from our MCMC fits. The lower dark curve and vertical dark line indicate the minimum specific heat and maximum neutrino luminosity compatible with \tilde{T} , respectively. The vertical gray region indicates the inferred L_ν . The horizontal gray regions indicate possible specific heat of the core. The curves marked $C = 2E/\tilde{T}$ and $L_\nu = L_{\text{in}}$, where L_{in} is the heating rate during an outburst, indicate limits on the specific heat and neutrino luminosity: The neutron star lies above and to the left of these curves. The inferred L_ν , for a uniform prior on Q_{imp} (blue histogram, Fig. 2), is indicated by the vertical gray band. The plot also shows the specific heat for scenarios in which the nucleons are unpaired throughout the core (upper gray band) and in which the nucleons are fully paired, so that only leptons contribute to the specific heat (lower gray band). The bands indicate the ranges of specific heat, computed using the models in Ref. [27]. The dashed lines indicate $\Delta T_{\text{eff}}^\infty/T_{\text{eff}}^\infty$ over a decade of cooling postoutburst.

the specific heat is $\lesssim 10^{37}$ erg K $^{-1}$ (lower horizontal gray band). Further modeling will be needed to separate the cooling of the pasta layer [30,31] from that of the core.

The direct Urca reactions require unpaired nucleons and a sufficiently large proton fraction. The proton fraction in neutron stars is controlled entirely by the poorly known density dependence of the symmetry energy, while the critical temperature for superfluidity and superconductivity is delicate and exponentially sensitive to the details of the assumed nuclear interactions. Requiring nuclear matter to be normal constrains the heat capacity of the core and provides a consistency check for nuclear models. Since the volume V_{dU} over which the direct Urca reaction operates is $\approx 1\%$ of the core, a large specific heat—indicated by $\Delta T_{\text{eff}}^\infty/T_{\text{eff}}^\infty \lesssim 1\%$ —would imply that the volume of the core with normal matter is $V_{\text{nrml}} \gg V_{\text{dU}}$. Having the onset pressure be $\approx 5\%$ below the central pressure of the neutron star would generate a sufficient V_{dU} to supply the required neutrino luminosity. Conversely, if $\Delta T_{\text{eff}}^\infty/T_{\text{eff}}^\infty \gtrsim 4\%$, then most of the neutrons in the star are in a superfluid phase, $V_{\text{nrml}} \approx V_{\text{dU}}$, and it is likely that the onset of direct Urca reactions is controlled by the closing of the neutron superfluid gap rather than by the proton fraction reaching the direct Urca threshold.

Assuming that fast cooling with $Q_f = 10^{36}$ erg km $^{-3}$ s $^{-1}$ turns on above a given threshold density (and hence mass), MXB 1659-29 is $\approx 0.03 M_\odot$ above the threshold mass, for typical equations of state. The transition may be broadened, however, over a range in density (see Ref. [34]); alternatively, the efficiency of the direct Urca reaction could be suppressed due to many-body effects. A value of $Q_f = 10^{35}$ erg km $^{-3}$ s $^{-1}$, expected if the nucleon effective masses were significantly reduced at a high density and if screening of the axial charge and two-body currents became important, would imply that the neutron star is not so close to the threshold mass. Proton pairing may persist with a critical temperature $T_c \gtrsim 10^7$ K over a significant volume in the core; the resulting suppression of the direct Urca rate by the factor $\simeq e^{-T_c/T}$ would favor a larger effective direct Urca volume.

Other reactions could also be responsible for the neutrino losses, for example, Urca reactions in the presence of a pion condensate or emission from unpaired quark matter. Both of these processes can cool the star rapidly, with a rate that scales as T^6 . The prefactor Q_f for these rates, although poorly known, is expected to be smaller, however, than direct Urca. If $Q_f \approx 10^{-1}$ of direct Urca, then these novel phases of matter would be compatible with observations of MXB 1659-29. If, on the other hand, $Q_f \approx 10^{-2}$ of direct Urca, pion condensation or unpaired quark matter are viable if most of the core were active, i.e., if the onset density for this cooling were close to saturation. Such a low onset density would imply, however, that MXB 1659-29 is significantly more massive than the observed slow-cooling

neutron stars. The neutrino luminosity of the transient neutron star KS 1731-260, for example, is $< 10^{-3}$ of the direct Urca luminosity [27]. The upper limits on thermal emission from SAX J1808.4-3658 [1,18,25,32,34] and 1H 1905 + 00 [17] imply, however, that neutron stars with much larger fast cooling volumes, and hence much higher central densities and masses, than MXB 1659-29 exist. We will explore the dependence on the equation of state in a future paper.

We have shown that, with three observed, regularly spaced accretion outbursts and a measured core temperature, the neutron star in MXB 1659-29 has a neutrino luminosity consistent with direct Urca reactions occurring in $\approx 1\%$ of the neutron star core. Taken together with KS 1731-260 [27], our work is further evidence that neutrino cooling in neutron star cores can be either fast or slow. Observations of the neutron star's effective temperature, now that the source has returned to quiescence, will measure the heat capacity of the core and help to untangle the physics underlying the neutrino luminosity. The fraction of unpaired normal particles implied by the heat capacity (Fig. 3) will distinguish whether the direct Urca threshold is achieved only towards the center of the star or superfluidity suppresses direct Urca throughout much of the core. This motivates an improved calculation of the direct Urca rate, which we shall address in a forthcoming publication.

We thank C. Heinke and W. Newton for useful discussion. This work benefited from discussions at the Physics and Astrophysics of Neutron Star Crusts workshop 2016 supported by the National Science Foundation under Grant No. PHY-1430152 (JINA Center for the Evolution of the Elements). A. C. is supported by an NSERC Discovery Grant and is a member of the Centre de Recherche en Astrophysique du Québec (CRAQ). E. F. B. is supported by the U.S. National Science Foundation Grant No. AST-1516969. F. J. F. and C. J. H. are supported in part by DOE Grants No. DE-FG02-87ER40365 and No. DE-SC0008808. D. P.'s work is partially supported by Conacyt through the Fondo Sectorial de Investigación para la Educación, Grant CB-2014-1, No. 240512. S. R. acknowledges support from the U.S. Department of Energy Grant No. DE-FG02-00ER41132. We thank the Aspen Center for Physics, which is supported by National Science Foundation Grant No. PHY-1607611, for a stimulating venue for completing this work.

*ebrown@pa.msu.edu

†andrew.cumming@mcgill.ca

‡ffattoye@indiana.edu

§horowitz@indiana.edu

||page@astro.unam.mx

¶sareddy@uw.edu

[1] D. G. Yakovlev and C. J. Pethick, *Annu. Rev. Astron. Astrophys.* **42**, 169 (2004).

- [2] D. Page, U. Geppert, and F. Weber, *Nucl. Phys.* **A777**, 497 (2006).
- [3] D. G. Yakovlev and P. Haensel, *Astron. Astrophys.* **407**, 259 (2003).
- [4] D. Page and S. Reddy, *Annu. Rev. Nucl. Part. Sci.* **56**, 327 (2006).
- [5] J. M. Lattimer, C. J. Pethick, M. Prakash, and P. Haensel, *Phys. Rev. Lett.* **66**, 2701 (1991).
- [6] O. Maxwell, G. E. Brown, D. K. Campbell, R. F. Dashen, and J. T. Manassah, *Astrophys. J.* **216**, 77 (1977).
- [7] G. E. Brown, K. Kubodera, D. Page, and P. Pizzochero, *Phys. Rev. D* **37**, 2042 (1988).
- [8] N. Iwamoto, *Phys. Rev. Lett.* **44**, 1637 (1980).
- [9] T. Schäfer and K. Schwenzer, *Phys. Rev. D* **70**, 114037 (2004).
- [10] E. Flowers, M. Ruderman, and P. Sutherland, *Astrophys. J.* **205**, 541 (1976).
- [11] D. N. Voskresenskii and A. V. Senatorov, *Zh. Eksp. Teor. Fiz.* **90**, 1505 (1986) [*JETP* **63**, 885 (1986)].
- [12] L. B. Leinson, *Phys. Rev. C* **84**, 045501 (2011).
- [13] H.-Y. Chiu and E. E. Salpeter, *Phys. Rev. Lett.* **12**, 413 (1964).
- [14] D. Viganò, N. Rea, J. A. Pons, R. Perna, D. N. Aguilera, and J. A. Miralles, *Mon. Not. R. Astron. Soc.* **434**, 123 (2013).
- [15] D. Page, J. M. Lattimer, M. Prakash, and A. W. Steiner, *Astrophys. J.* **707**, 1131 (2009).
- [16] M. Colpi, U. Geppert, D. Page, and A. Possenti, *Astrophys. J. Lett.* **548**, L175 (2001).
- [17] P. G. Jonker, D. Steeghs, D. Chakrabarty, and A. M. Juett, *Astrophys. J. Lett.* **665**, L147 (2007).
- [18] C. O. Heinke, P. G. Jonker, R. Wijnands, C. J. Deloye, and R. E. Taam, *Astrophys. J.* **691**, 1035 (2009).
- [19] R. Wijnands, M. P. Muno, J. M. Miller, L. M. Franco, T. Strohmayer, D. Galloway, and D. Chakrabarty, *Astrophys. J.* **566**, 1060 (2002).
- [20] R. Wijnands, M. Nowak, J. M. Miller, J. Homan, S. Wachter, and W. H. G. Lewin, *Astrophys. J.* **594**, 952 (2003).
- [21] E. M. Cackett, R. Wijnands, M. Linares, J. M. Miller, J. Homan, and W. H. G. Lewin, *Mon. Not. R. Astron. Soc.* **372**, 479 (2006).
- [22] E. M. Cackett, R. Wijnands, J. M. Miller, E. F. Brown, and N. Degenaar, *Astrophys. J.* **687**, L87 (2008).
- [23] E. M. Cackett, E. F. Brown, A. Cumming, N. Degenaar, J. K. Fridriksson, J. Homan, J. M. Miller, and R. Wijnands, *Astrophys. J.* **774**, 131 (2013).
- [24] C. Sanchez-Fernandez, D. Eckert, E. Bozzo, J. Kajava, E. Kuulkers, and J. Chenevez, *Astronomer's Telegram* **7946**, 1 (2015).
- [25] R. Wijnands, N. Degenaar, and D. Page, *Mon. Not. R. Astron. Soc.* **432**, 2366 (2013).
- [26] P. Haensel and J. L. Zdunik, *Astron. Astrophys.* **480**, 459 (2008).
- [27] A. Cumming, E. F. Brown, F. J. Fattoyev, C. J. Horowitz, D. Page, and S. Reddy, *Phys. Rev. C* **95**, 025806 (2017).
- [28] E. M. Cackett, E. F. Brown, A. Cumming, N. Degenaar, J. M. Miller, and R. Wijnands, *Astrophys. J. Lett.* **722**, L137 (2010).

- [29] E. F. Brown and A. Cumming, *Astrophys. J.* **698**, 1020 (2009).
- [30] C. J. Horowitz, D. K. Berry, C. M. Briggs, M. E. Caplan, A. Cumming, and A. S. Schneider, *Phys. Rev. Lett.* **114**, 031102 (2015).
- [31] A. Deibel, A. Cumming, E. F. Brown, and S. Reddy, *Astrophys. J.* **839**, 95 (2017).
- [32] C. O. Heinke, P. G. Jonker, R. Wijnands, and R. E. Taam, *Astrophys. J.* **660**, 1424 (2007).
- [33] D. K. Galloway, M. P. Muno, J. M. Hartman, D. Psaltis, and D. Chakrabarty, *Astrophys. J. Suppl. Ser.* **179**, 360 (2008).
- [34] M. V. Beznogov and D. G. Yakovlev, *Mon. Not. R. Astron. Soc.* **452**, 540 (2015).